



Aquasteril

- ◆ Removes slime deposits (biofilm)
- ◆ Removes deposits such as slimy vitamin and drug residues
- ◆ Is effective against bacteria, fungi and algae
- ◆ Is suitable for organic farming
- ◆ Biodegradable

Used to sanitize and clean drinking water, water lines, water tanks, drinkers, and nipple drinkers, particularly in poultry and pig housing.

ACTIVE INGREDIENTS: 50mg/g peracetic acid, 265 mg/g hydrogen peroxide

Drinking water sanitation during the fattening process

Aquasteril	+ water	= ready-to-use solution
20ml	1000 litres	0.002%

Depending on the microorganism, add Aquasteril to the drinking water over a period of approx. 14 days.

Sanitizing and cleaning water lines in empty animal housing

Aquasteril	+ water	= ready-to-use solution
2 litres	100 litres	2%

Listed by FiBL

Before the animals enter the housing, pump Aquasteril into the drinking water system. Ensure sufficient ventilation. Open all nipple and the ends of the water lines (wear protective clothing). The rising air bubbles indicate whether Aquasteril has reached all parts of the water system. Allow the solution either to circulate for six hours or close the system for 12 to 24 hours. Then, rinse the whole system with clean water. Close all nipples and drinking water line valves.

Aquasteril can be used in ECOLOGICAL farms

Water disinfectants and the use of water soluble "blue dye":

In order to optimise the administration of Aquasteril at the end-of-cycle it is useful to add an approved "blue dye" to the disinfectant. This will help ensure that the system is initially filled with disinfectant, and later show that the system is fully drained.

Disinfectants and Biological products:

If live bacterial vaccines or probiotics are to be administered orally to the flock, it may be necessary to withdraw Aquasteril 3 days prior to their administration. In addition Aquasteril should only be re-introduced into the drinking water 3 days post vaccination.

Corrosion data and user precautions:

!! Please pay careful attention to the following information: - Peracetic acid and Hydrogen Peroxide are corrosive and should only be administered through systems fitted with Teflon pumps (alternatively pre-dilute to a 1% solution before the solution enters the pump) - Never mix Aquasteril with other products! - A safety data sheet is available upon request.

General protective and hygienic measures:

When using/working with concentrated disinfectants do not eat, drink or smoke. Avoid contact with the undiluted product. Wash hands after work.

**Hazard warnings for the concentrate:**

Marked with the pictograms GHS02, GHS05, GHS07, GHS09 / Signal word: Hazard
H statements: H242, H290, H302, H312, H314, H332, H335, H410

baua reg. No.: N-55850 (PT 3), N-55851 (PT 4), N-55852 (PT 5)

Breathing protection	For poorly ventilated areas - filter apparatus type A
Hand protection	Protective rubber gloves e.g. Nitril
Eye protection	Glasses with side protection, or Basket eye glasses

Wash hands after work.

Pack size:
10kg

Use Biocides Safely.
Always read the label and product information before use!