



Interkokask Concentrate

- ◆ Bacteria, Viruses and Fungi
- ◆ Bacterial Spores
- ◆ Mycobacterium (TB)
- ◆ Coccidial Oocysts and Cryptosporidium
- ◆ Breaks through biofilm
- ◆ BEIC Approved

Interkokask Concentrate has a special formula including chlorocresol which is a lipid-solvent that breaks through the protective triple outer protein/lipid/chitin layers of worm eggs and coccidial oocysts. **Interkokask Concentrate** removes immature parasites, which means that only one application is sufficient to remove them from the environment.

ACTIVE INGREDIENTS: Chlorocresol.

UK DEFRA Approvals:

Poultry Orders	2.0%	30 minutes
Tuberculosis Orders	3.0%	60 minutes
Foot and Mouth Disease	4.0%	30 minutes
Swine Vesicular Disease	4.0%	30 minutes

DVG (German Veterinary Association) Approvals:

Disinfection of coccidial oocysts	4.0%	2 hours
Disinfection of cryptosporidium	2.0%	2 hours
Disinfection of worms and worm eggs	2.0%	2 hours (non-UK)

FAO Resistance Test:

Disinfection of red mite and red mite eggs	4.0%	24 hours (non EU)
--	-------------	-------------------

European approvals:

EN 14349/14675 Bacteria/Virus:

Bacteria (light soiling)	0.50%	30 minutes
Bacteria (heavy soiling)	0.50%	30 minutes
Viruses	4.00%	30 minutes

EN 1657 Fungi:

Fungi (light soiling)	2.00%	30 minutes
EN 13704 Spores:		
Bacterial spores	4.00%	60 minutes

EN 14204 Mycobacterium:

Mycobacterium (Light soiling)	0.50%	60 minutes
Mycobacterium (Heavy soiling)	1.00%	60 minutes

Ultra wide spectrum disinfectant!

Treating	Minimum exposure	Dilution
Red mite / Red mite eggs	24 hours - treat whole animal house	3% (300ml in 10 litres of water)
Coccidial oocysts / Cryptosporidia	2 hours - treat floors and walls up to 2 metres	4% (400ml in 10 litres of water)
Worms / Worm eggs	2 hours - treat floors and walls up to 2 metres	2% (200ml in 10 litres of water)
Mycobacterium	1 hour - treat whole animal house	1% (100ml in 10 litres of water)
Bacterial spores	1 hour - treat whole animal house	4% (400ml in 10 litres of water)
Bacteria	30 minutes - treat whole animal house	1% (100ml in 10 litres of water)
Fungi	30 minutes - treat whole animal house	3% (300ml in 10 litres of water)
Viruses	30 minutes - treat whole animal house	3% (300ml in 10 litres of water)
Unknown Diseases (except red mite)	2 hours - treat whole animal house	4% (400ml in 10 litres of water)

Important: Ideally, before and after use, rinse all tools and plastic parts (e.g. high pressure tools, hose pipes) thoroughly with a diluted alkaline detergent solution. Any traces of formaldehyde, glutaraldehyde, or rust may cause the chlorocresol to crystallise. Do not pre-dilute the product prior to use.

Note: Interclean® Foam Cleaner is specially formulated to clean equipment used for spraying **Interkokask** and **Interkokask Concentrate**. For more information, please contact Hysolv Ltd



INSTRUCTIONS FOR USE

- May be used on vehicles, machinery and equipment:** not known to be corrosive when used at recommended dilutions.
- Works at low temperatures:** DEFRA tested at 4°C.
- Disinfects in presence of organic matter:** tested on absorbent surfaces and in presence of organic matter.
- Footbaths:** Bacteria/viruses/fungi (coccidial oocysts) – **2.00% (4.00%)**, change solution weekly or more often if heavily soiled.



- ◆ Remove animals and clean the housing thoroughly.
- ◆ Rinse all spray and mixing equipment thoroughly with water prior to use.
- ◆ Make sure that all traces of previously used chemicals are removed.
- ◆ Adjust the sprayer to deliver the required dilution of Interkokask Concentrate. Alternatively fill the sprayer tank with clean water, allowing enough room for the Interkokask Concentrate to be added. (Please note, on cold days add warm water to the tank to bring the temperature to 7°C – 15°C.)
- ◆ Add Interkokask Concentrate into the sprayer's tank (bucket if using a knapsack sprayer) and mix thoroughly to make the required concentration of product.
- ◆ Apply the diluted product on the dry surfaces of the empty animal house with suitable spraying equipment.
- ◆ Coccidia/Cryptosporidium: Treat the floors and walls up to a minimum of 2 metres. Walls above 2 metres and ceilings should be treated only if advised. Spray all equipment and built-in parts. Viruses/Bacteria/Fungi/Mycobacteria: Treat the whole house.
- ◆ The buildings and equipment should be treated for 2 – 24 hrs as shown in the 'treatment times' chart overleaf.
- ◆ After the building is dry, re-equip and re-stock

Spray Application

- ◆ High pressure spraying equipment: Adjust down to a max of 1000 psi (70 bar), or if possible 300 psi (20 bar) to reduce spray "bounce-back" during application to walls. Use a "wide angle fan spray nozzle"
 - ◆ Knapsack spraying equipment: Use a "wide angle fan spray nozzle"
- Apply a minimum of 100ml solution per square metre on concrete and other smooth surfaces, Wooden or porous surfaces should be saturated and may require up to 400ml solution / m².

1 kg Interkokask Concentrate

- ◆ . 0.50% maximum treatable area approx. 1,920 m² - 480 m² *
 - ◆ . 1.00% maximum treatable area approx. 960 m² - 240 m² *
 - ◆ . 2.00% maximum treatable area approx. 480 m² - 120 m² *
 - ◆ . 3.00% maximum treatable area approx. 320 m² - 80 m² *
 - ◆ . 4.00% maximum treatable area approx. 240 m² - 60 m² *
- * = depending on application rate.

When using/working with concentrated disinfectants do not eat, drink or smoke. Avoid contact with eyes skin and clothing. Wear the recommended protective clothing and equipment - see below: Wash hands after work.

Breathing protection	Full face mask, filter A2 B2-P2
Hand protection	Protective rubber gloves e.g. Nitril
Eye protection	Full mask
Body protection	Chemical-protective clothing

Pack sizes:

1kg, 10kg, 1000kg

Use Biocides Safely.

Always read the label and product information before use!

Product disposal: In most instances, Interkokask Concentrate diluted to approved concentrations may be disposed of through the normal drainage systems. Please check with local authorities if unsure of regional waste disposal requirements.